

WizPure™ Bst DNA Polymerase (Full Length)

RUO For Research Use Only

REF W1320

DESCRIPTION

WizPure™ Bst DNA Polymerase (Full Length) is the full length polymerase from *Bacillus stearothermophilus* that contains the 5' → 3' polymerase activity and the 5' → 3' exonuclease activity, but lacks 3' → 5' exonuclease activity.

KIT CONTENTS

Contents	W1320	W1320-5
Bst DNA Polymerase (FL), (8U/μl)	250 μl	1,250 μl
10X Bst Reaction Buffer	1 ml	5 x 1 ml
100mM MgSO ₄	1 ml	5 x 1 ml

STORAGE BUFFER

10mM Tris-HCl, 50mM KCl, 1.0mM dithiothreitol, 0.1mM EDTA, 0.1% Triton X-100, 50% glycerol, pH 7.5 (25°C)

10X Bst REACTION BUFFER

200mM Tris-HCl, 100mM Ammonium Sulfate, 100mM KCl, 1.0% Triton X-100, pH 8.8 (25°C)

ACTIVITY

- 5'→3' exonuclease: Yes (double-strand specific)
- 3'→5' exonuclease: No
- Strand Displacement: No

APPLICATIONS

- Isothermal DNA amplification
- Primer extension
- Nick translation

UNIT DEFINITION

1 unit is defined as the amount of polymerase required to convert 10 nmol of dNTPs into acid insoluble material in 30 minutes at 65°C.

STORAGE CONDITIONS

Store all components at -20°C in a non-frost-free freezer.

QUALITY CONTROL ANALYSIS:

In accordance with Wizbiosolutions Inc. ISO 13485-certified Quality Management System, each lot of WizPure™ Bst DNA Polymerase (Full Length) is tested against predetermined specifications to ensure consistent product quality.

ORDERING INFORMATION

Product	Cat No.	Package
WizPure™ Bst DNA Polymerase (Full Length)	W1320	2,000 U
	W1320-5	10,000 U

PROTOCOL

1. LAMP Reaction Mixture Preparation

Prior to the experiment, it is prudent to carefully optimize experiment conditions and to include controls at every stage. See pre-protocol considerations for details.

For multiple reactions, scale up the volume of reaction components proportionally. All reagents can be prepared at room temperature.

1. Assemble reaction tubes on ice whenever possible to avoid premature, nonspecific polymerase activity.
2. The following table shows recommended component volumes:

Component	25 μl reaction	Final Conc.
Bst DNA Polymerase (FL)	1 - 2 μl	
10X Bst Reaction Buffer	2.5 μl	1X
dNTP mix (each 10mM)	3.5 μl	1.4 mM
100mM MgSO ₄	2 - 2.5 μl	8 - 10 mM
FIP Primer (10 μM)	2 - 5 μl	0.8 - 2.0 μM
BIP primer (10 μM)	2 - 5 μl	0.8 - 2.0 μM
F3 primer (10 μM)	0.5 - 1 μl	0.2 - 0.4 μM
R3 primer (10 μM)	0.5 - 1 μl	0.2 - 0.4 μM
Loop F (10 μM)	1 - 2 μl	0.4 - 0.8 μM
Loop B (10 μM)	1 - 2 μl	0.4 - 0.8 μM
Green fluorescence dye (25X)*	1 μl	1X
Template DNA	1 - 5 μl	
Distilled water	up to 25 μl	

* For Real-time fluorescence assay only

3. Ensure reactions are mixed thoroughly by pipetting or gentle vortexing followed by a brief spin in a microcentrifuge.

For Conventional LAMP Assay

1. Transfer tubes on ice into an isothermal instrument or water bath.
2. Incubation 62°C for 30-60 min.

NOTE: Assay conditions may need to be optimized, depending on different primer and template combinations.

3. Run 5μ of the amplification product on a 1.5% agarose gel.

For Conventional LAMP Assay

1. Transfer tubes to a Real-time assay instrument (eg, Real-time PCR machine)
2. Real-time LAMP program setting and running as follows,
 - Incubation temperature: 62°C
 - Time: 30-60 min (Scan/1 min).
 - Detection channel: SYBR green (FAM)

Technical Support



www.wizbiosolution.com
support@wizbiosolution.com
+82 70 7605 5066



Wizbiosolutions Inc.
A-802, Woolim Lions Valley 2
Sagimakgol-ro 45beon-gil 14
Jungwon-gu, Seongnam-si, Gyeonggi-do,
13209 Republic of Korea

ISO13485:2016, GMP certified
WS-IFU-W1320-01(EU) (V.1.0.2, 20-10-23)